

Lecture slides (AI3501/CT4201/EC4215 – Computer Graphics)

# Culling

---

Lecturer: Bochang Moon

# Culling


---

- An optimization process that removes invisible geometry to speed up rendering
  
- Three types of culling
  - View volume culling
  - Occlusion culling
  - Back-face culling

# View Volume Culling


---

- A process to remove geometry that is outside the view volume
- Q. why do we need to do this culling?
- Q. how do we efficiently identify the object that is totally outside of the volume?



# View Volume Culling


- A process to remove geometry that is outside the view volume
- Q. why do we need to do this culling?



# View Volume Culling

---

- A process to remove geometry that is outside the view volume
- Q. how do we efficiently identify the object that is totally outside of the volume?
  - A bounding volume can be utilized. Why?



# View Volume Culling




---

- Simple bounding volumes

- Bounding box

- e.g., axis-aligned bounding box (AABB)


- Bounding sphere




# View Volume Culling

---


- Need identify the three cases



inside



intermediate




outside

# Background: Implicit Functions

---

- 2D implicit curves



- 3D implicit surfaces

- $f(x, y, z) = 0$

# Background: Implicit Functions

---

- Infinite plane through point  $\mathbf{a}$  with surface normal  $\mathbf{n}$ 
  - $(\mathbf{p} - \mathbf{a}) \cdot \mathbf{n} = 0$
  - The surface normal  $\mathbf{n}$  is a vector perpendicular to the plane.
  - When a point  $\mathbf{p}$  is on the plane,  $(\mathbf{p} - \mathbf{a}) \cdot \mathbf{n}$  will be zero.
    - Recall the definition of a dot product
      - $\mathbf{a} \cdot \mathbf{b} = \|\mathbf{a}\| \|\mathbf{b}\| \cos\theta$

# View Volume Culling



- We can check the following:

- $\frac{(c-a) \cdot n}{\|n\|} > r$

- **c**: center of the bounding sphere


- r: radius of the sphere

- Q. what's the geometric meaning of  $\frac{(c-a) \cdot n}{\|n\|}$ ?




# Background: Dot Product

- Vector multiplications
  - Dot product (scalar product)
    - $\mathbf{a} \cdot \mathbf{b} = \|\mathbf{a}\| \|\mathbf{b}\| \cos\theta$
    - Usage: ( $\mathbf{a} \rightarrow \mathbf{b}$ ) projection of a vector to another one
    - $\mathbf{a} \rightarrow \mathbf{b} = \|\mathbf{a}\| \cos\theta = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{b}\|}$
    - Note: this is the length of the projected vector onto  $\mathbf{b}$
  - Dot product in Cartesian coordinates
    - Properties:  $\mathbf{x} \cdot \mathbf{x} = \mathbf{y} \cdot \mathbf{y} = 1$  and  $\mathbf{x} \cdot \mathbf{y} = 0$
    - $\mathbf{a} \cdot \mathbf{b} = (x_a \mathbf{x} + y_a \mathbf{y}) \cdot (x_b \mathbf{x} + y_b \mathbf{y})$
    - $= x_a x_b (\mathbf{x} \cdot \mathbf{x}) + x_a y_b (\mathbf{x} \cdot \mathbf{y}) + x_b y_a (\mathbf{y} \cdot \mathbf{x}) + y_a y_b (\mathbf{y} \cdot \mathbf{y})$
    - $= x_a x_b + y_a y_b$
    - In 3D,
      - $\mathbf{a} \cdot \mathbf{b} = x_a x_b + y_a y_b + z_a z_b$




# View Volume Culling

- Need identify the three cases




inside

$$\frac{(\mathbf{c} - \mathbf{a}) \cdot \mathbf{n}}{\|\mathbf{n}\|} < -r$$



intermediate

$$-r < \frac{(\mathbf{c} - \mathbf{a}) \cdot \mathbf{n}}{\|\mathbf{n}\|} < r$$




outside

$$\frac{(\mathbf{c} - \mathbf{a}) \cdot \mathbf{n}}{\|\mathbf{n}\|} > r$$

# View Volume Culling

---

- Q. can we optimize our pipeline further?





intermediate

$$-r < \frac{(\mathbf{c} - \mathbf{a}) \cdot \mathbf{n}}{\|\mathbf{n}\|} < r$$

# Hierarchical Culling



- If a bounding volume is intermediate,
  - Check its left and right children



# Back-Face Culling

- If the angle between the view and normal is within a range (-90 to 90 degrees), the triangle is visible.

o  $\cos \theta \geq 0$



# Back-Face Culling



- If the angle between the view and normal is within a range (-90 to 90 degrees), the triangle is visible.

- $\cos\theta \geq 0$

- $e \cdot n \geq 0$

- Dot product


- $a \cdot b = \|a\| \|b\| \cos\theta$



# Back-Face Culling

---

- Assumption for the back-face culling:
  - Models are closed (i.e., no holes).



# Further Readings

---

- Chapter 2.5
- Chapter 8.4 and 12